arXiv:1209.3006v1 [math.PR] 13 Sep 2012

A generalized telegraph process with velocity driven by random trials

Irene Crimaldi[*], Antonio Di Crescenzo, Antonella Iuliano, Barbara Martinucci[†]

September 14, 2012

Abstract

We consider a random trial-based telegraph process, which describes a motion on the real line with two constant velocities along opposite directions. At each epoch of the underlying counting process the new velocity is determined by the outcome of a random trial. Two schemes are taken into account: Bernoulli trials and classical Pólya urn trials. We investigate the probability law of the process and the mean of the velocity of the moving particle. We finally discuss two cases of interest: (i) the case of Bernoulli trials and intertimes having exponential distributions with linear rates (in which, interestingly, the process exhibits a logistic stationary density with non-zero mean), and (ii) the case of Pólya trials and intertimes having first Gamma and then exponential distributions with constant rates.

Keywords Telegraph process; random intertimes; Bernoulli scheme; Pólya urn model; logistic stationary density

Mathematics Subject Classification Primary 60K15; Secondary 60K37.

Running title Telegraph process driven by random trials.

1 Introduction

Since the 1950s several authors investigated the (integrated) telegraph process as a realistic model of random motion (see Goldstein [14], Bartlett [3] and Kac [17]). Such process describes a motion of a particle on the real line characterized by constant speed, the direction being reversed at the random epochs of a Poisson process. The probability density of the particle's position satisfies a hyperbolic differential equation, whose probabilistic properties have been studied for instance by Orsingher [20], [21], Foong and Kanno [13], and more recently by Beghin

---

[*] IMT Institute for Advanced Studies, Piazza San Ponziano 6, I-55100 Lucca, Italy, irene.crimaldi@imtlucca.it

[†] Dipartimento di Matematica, Università di Salerno, Via Ponte don Melillo, Fisciano (SA), I-84084, Italy, {adicrescenzo, aiuliano, bmartinucci}@unisa.it

et al. [4]. See also the book of Pinsky [22], in which many results on the telegraph process and its generalizations were given, and many applications were discussed. Certain one-dimensional generalizations of the telegraph process focus on cases in which the intertimes between two consecutive changes of direction are characterized by a variety of distributions. We recall the case of Erlang distribution (Di Crescenzo [9]), of gamma distribution (Di Crescenzo and Martinucci [10]), of the exponential distribution with linearly increasing rate (Di Crescenzo and Martinucci [11]). Moreover, investigations on the telegraph process with random velocities have been performed by Stadje and Zacks [24], and by De Gregorio [8], whereas a generalized telegraph process governed by an alternating renewal process was studied by Zacks [26]. Iacus [16] gave a rare example where an explicit probability law is obtained for an inhomogeneous telegraph process.

In this paper we aim to study the telegraph model subject to a further source of randomness, by assuming that the velocity of the moving particle is driven by random trials. Precisely, we deal with a two-velocity random motion on the real line where, differently from the classical telegraph process whose positive and negative velocities are alternating, at each time epoch now the new velocity is determined by the outcome of a random trial. The latter follows one out of two schemes: the *Bernoulli scheme*, which acts on independence, and the classical *Pólya scheme* (cf. Pólya [23], Mahmoud [18]), where the outcome of each trial depends on the outcomes of the previous ones.

The paper is organized as follows. In Section 2 we describe in detail the mathematical model of the motion. In Section 3, we investigate the probability law and the conditional mean velocity of the process. Then, in Section 4, for the Bernoulli scheme we discuss the instance in which the random intertimes between consecutive trials are exponentially distributed with linearly increasing rates. In this case we obtain the probability density of the process in closed form, we show that it possesses a logistic stationary density, and then express the conditional mean velocity of the process in terms of hypergeometric functions. Finally, in Section 5, we show a case for the Pólya scheme in which the first random intertimes in both directions are Gamma-distributed, whereas all remaining intertimes are exponentially distributed. We obtain the probability density in closed form and the conditional mean velocity as series of Gauss hypergeometric functions. For the reader's convenience, the paper is enriched by two Appendices containing definitions and formulas used in the proofs.

2 Stochastic model

Let $\{(S_t, V_t); t \geq 0\}$ be a continuous-time stochastic process, where S_t and V_t denote respectively position and velocity at time t of the moving particle. The motion is characterized by two velocities, c and $-v$, with $c, v > 0$, the direction of the motion being specified by the sign of the velocity. At time $T_0 = 0$ the particle starts from the origin, thus $S_0 = 0$. The initial velocity V_0 is determined by the outcome X_1 of the first random trial. At the random time $T_1 > 0$ the particle is subject to an event, whose effect possibly changes the velocity according to the outcome of the second random trial. This behavior is repeated cyclically at every instant of a sequence of random epochs $T_0 = 0 < T_1 < T_2 < T_3 < \cdots$. We assume that the durations of time intervals $[T_n, T_{n+1})$, $n = 0, 1, 2, \ldots$, constitute a sequence of non-negative random variables. Precisely, let U_k (resp. D_k) denote the random duration of the k-th time period during which the particle moves forward, with velocity c (resp. backward, with velocity $-v$). Furthermore, $\{U_k; k \geq 1\}$ and $\{D_k; k \geq 1\}$ are mutually independent sequences of non-negative and absolutely continuous independent random variables. Denoting by Z_n the velocity of the particle during the interval $[T_n, T_{n+1})$, we assume that $\{Z_n; n \geq 0\}$ is a sequence of random variables governed by the sequence $\{X_n; n \geq 1\}$ of the outcomes of the random trials. Moreover we assume that the collection $\{U_k, D_k; k \geq 1\}$ is independent of $\{X_n; n \geq 1\}$.

In this paper we focus on the case in which the random trials $\{X_n; n \geq 1\}$ are the outcomes of a sequence of indicator functions such that

$$\mathrm{P}\{Z_0 = c\} = \mathrm{P}\{X_1 = 1\} = \frac{b}{b+r}, \qquad \mathrm{P}\{Z_0 = -v\} = \mathrm{P}\{X_1 = 0\} = \frac{r}{b+r}. \tag{1}$$

and for $n \geq 1$

$$\begin{aligned} \mathrm{P}\{Z_n = c \,|\, \mathcal{G}_n\} = \mathrm{P}\{X_{n+1} = 1 \,|\, \mathcal{G}_n\} &= \frac{b + A\sum_{k=1}^{n} X_k}{b+r+An}, \\ \mathrm{P}\{Z_n = -v \,|\, \mathcal{G}_n\} = \mathrm{P}\{X_{n+1} = 0 \,|\, \mathcal{G}_n\} &= \frac{r + A\sum_{k=1}^{n}(1 - X_k)}{b+r+An}, \end{aligned} \tag{2}$$

where b and r are positive constants, A is a non-negative constant, $\mathcal{G}_0 = \{\emptyset, \Omega\}$ and $\mathcal{G}_n = \sigma(X_1, \ldots, X_n)$ for $n \geq 1$.

We can distinguish two different cases.

1) The case $A = 0$, which means that the random trials $\{X_n; n \geq 1\}$ are independent, i.e. they are a *Bernoulli scheme* with parameter

$$p = \frac{b}{b+r}. \tag{3}$$

2) The case $A > 0$, which means that $\{X_n; n \geq 1\}$ are the outcomes of a sequence of drawings from an urn which initially has b black balls and r red balls and then is updated according to the *classical Pólya urn scheme*: the drawn ball is returned into the urn together with $A > 0$ balls of the same color.[1] ($X_1 = 1$ means that the drawn ball is black; otherwise we have $X_1 = 0$.)

   In Section 5.1 we discuss an extension of the above setting to the case in which $\{X_n; n \geq 1\}$ is the sequence of the outcomes of drawings from a *randomly reinforced urn*.

The novelty of the present model of random motion is the inclusion of a sequence of random trials that regulates the velocity of the particle.

---

[1] We recall that, from a mathematical point of view, also in this case the parameters b, r, A of the model can be *real*.



Figure 1: Sample paths of S_t with *(a)* $V_0 = c$, and *(b)* $V_0 = -v$.

Let M_t be the stochastic process which counts the number of epochs T_i (with $i \geq 1$) occurring before t, i.e.

$$M_t = \max\{i \geq 1 : T_i \leq t\}, \qquad t > 0. \tag{4}$$

It is worthwhile to note that position and velocity of the particle at time t can thus be formally expressed as:

$$V_t = Z_{M_t}, \qquad S_t = \int_0^t V_s \,\mathrm{d}s, \qquad t > 0. \tag{5}$$

Figure 1 shows two examples of sample paths of S_t, with indication of the random intertimes U_k and D_k, where the sequence $\{X_n; n \geq 1\}$ takes values *(a)* $\{1, 1, 0, 0, 1, 0, 1, \ldots\}$ and *(b)* $\{0, 1, 1, 0, 1, 1, \ldots\}$, respectively.

The following stochastic equation holds:

$$S_{T_{k+1}} = S_{T_k} + W_k, \qquad k \geq 0\,,$$

where $\{W_k; k \geq 0\}$ is the sequence of random variables defined by

$$W_0 = \begin{cases} cU_1 & \text{if } Z_0 = c \\ -vD_1 & \text{if } Z_0 = -v, \end{cases}$$

and, for $k \geq 1$ and $1 \leq j \leq k+1$,

$$W_k = \begin{cases} cU_j & \text{if } Z_k = c \text{ and } X_1 + N_{k-1} = j - 1 \\ -vD_j & \text{if } Z_k = -v \text{ and } X_1 + N_{k-1} = k - j + 1\,, \end{cases}$$

where N_{k-1} is the random variable that counts the number of random trials yielding velocity c among the trials going from the 2-nd to the k-th one, i.e.

$$N_{k-1} = \sum_{i=1}^{k-1} \mathbf{1}_{\{Z_i = c\}} = \sum_{i=1}^{k-1} X_{i+1} = \sum_{h=2}^{k} X_h\,, \qquad k \geq 2. \tag{6}$$

For convenience, we set also $N_0 = 0$. In Appendix A we give some useful formulas regarding the probability distribution of N_{k-1} conditioned on the initial velocity for the two different schemes.

Throughout the paper we denote by f_{U_k} (resp. F_{U_k}, $\overline{F}_{U_k}$) and by f_{D_k} (resp. F_{D_k}, $\overline{F}_{D_k}$) the probability densities (resp. cumulative distribution functions, tail distribution functions) of U_k and D_k, respectively. Moreover, $f_U^{(k)}$ and $f_D^{(k)}$ (resp., $F_U^{(k)}$ and $F_D^{(k)}$, $\overline{F}_U^{(k)}$ and $\overline{F}_D^{(k)}$) will denote the probability densities (resp., cumulative distribution functions, tail distribution functions) of the partial sums

$$U^{(k)} = U_1 + U_2 + \cdots + U_k, \qquad D^{(k)} = D_1 + D_2 + \cdots + D_k, \qquad k \geq 1\,.$$

3 Probability law and mean velocity

At time $t = 0$ we assume $S_0 = 0$ so that at time $t > 0$ the particle is located in the domain $[-vt, ct]$. The probability law of S_t, $t > 0$, thus possesses a discrete component on the points $-vt$ and ct, and an absolutely continuous component over $(-vt, ct)$, which will be investigated in the sequel.

Proposition 3.1 *For all $t > 0$ we have*

$$\mathrm{P}\{S_t = ct\} = \frac{b}{b+r}\overline{F}_{U_1}(t) + \sum_{k=1}^{+\infty} \mathrm{P}\{Z_0 = c, N_k = k\}\left[F_U^{(k)}(t) - F_U^{(k+1)}(t)\right], \tag{7}$$

where

$$\mathrm{P}\{Z_0 = c, N_k = k\} = \begin{cases} p^{k+1} & \text{if } A = 0 \\ \dfrac{b}{b+r}\left(\dfrac{b+A}{A}\right)_k \left[\left(\dfrac{b+A+r}{A}\right)_k\right]^{-1} & \text{if } A > 0. \end{cases}$$

Similarly, we have

$$\mathrm{P}\{S_t = -vt\} = \frac{r}{b+r}\overline{F}_{D_1}(t) + \sum_{k=1}^{+\infty} \mathrm{P}\{Z_0 = -v, N_k = 0\}\left[F_D^{(k)}(t) - F_D^{(k+1)}(t)\right], \tag{8}$$

where

$$\mathrm{P}\{Z_0 = -v, N_k = 0\} = \begin{cases} (1-p)^{k+1} & \text{if } A = 0 \\ \dfrac{r}{b+r}\left(\dfrac{r+A}{A}\right)_k \left[\left(\dfrac{b+A+r}{A}\right)_k\right]^{-1} & \text{if } A > 0. \end{cases}$$

Proof. First of all, note that, for $y \in \{-v, c\}$, condition $S_t = yt$ implies $V_0 = y$ and $V_t = y$ a.s. Conditioning on $M_t = k$ and recalling (6) we have

$$\begin{aligned} \mathrm{P}\{S_t = ct\} &= \mathrm{P}\{S_t = ct, V_t = c, V_0 = c\} \\ &= \sum_{k=0}^{+\infty} \mathrm{P}\{Z_0 = c, N_k = k\}\, \mathrm{P}\{M_t = k \,|\, Z_0 = c, N_k = k\}. \end{aligned}$$

Eq. (7) thus easily follows from (1) and from Eqs. (45) and (50) for $j = k - 1$. In the same way Eq. (8) follows from (1) and from Eqs. (48) and (53). □

Let us now define the probability density of S_t, $t > 0$, conditional on initial velocity $y \in \{-v, c\}$:

$$p(x,t\,|\,y) = \frac{\partial}{\partial x}\mathrm{P}\{S_t \le x\,|\,V_0 = y\}, \qquad x \in (-vt, ct). \tag{9}$$

We remark that, for $t > 0$,

$$\mathrm{P}\{S_t = yt\,|\,V_0 = y\} + \int_{-vt}^{ct} p(x,t\,|\,y)\,\mathrm{d}x = 1, \qquad y \in \{-v, c\}.$$

The density of the particle position is

$$p(x,t) = \frac{\partial}{\partial x}\mathrm{P}\{S_t \le x\} = \frac{b}{b+r}\,p(x,t\,|\,c) + \frac{r}{b+r}\,p(x,t\,|\,-v), \tag{10}$$

where $p(x,t\,|\,y)$, defined in (9), can be expressed as

$$p(x,t\,|\,y) = f(x,t\,|\,y) + b(x,t\,|\,y). \tag{11}$$

Here f and b denote the densities of particle's position when the motion at time t is characterized by forward and backward velocity, respectively, i.e.

$$\begin{aligned} f(x,t\,|\,y) &= \frac{\partial}{\partial x}\mathrm{P}\{S_t \le x, V_t = c\,|\,V_0 = y\}, \\ b(x,t\,|\,y) &= \frac{\partial}{\partial x}\mathrm{P}\{S_t \le x, V_t = -v\,|\,V_0 = y\}, \end{aligned}$$

for x in $(-vt, ct)$. Then, for $y \in \{-v, c\}$, we can write

$$f(x,t\,|\,y) = \sum_{k=1}^{+\infty} f_k(x,t\,|\,y), \qquad b(x,t\,|\,y) = \sum_{k=1}^{+\infty} b_k(x,t\,|\,y) \tag{12}$$

where, for x in $(-vt, ct)$,

$$f_k(x,t\,|\,y) = \frac{\partial}{\partial x}\mathrm{P}\{S_t \le x, V_t = c, M_t = k\,|\,V_0 = y\}, \tag{13}$$

$$b_k(x,t\,|\,y) = \frac{\partial}{\partial x}\mathrm{P}\{S_t \le x, V_t = -v, M_t = k\,|\,V_0 = y\}, \tag{14}$$

with M_t defined in (4). Hereafter we give an expression for densities f_k and b_k conditioned by $V_0 = c$. Note that $f_1(x,t\,|\,c) = 0$.

Theorem 3.1 *For $t > 0$ and $-vt < x < ct$, the densities (13) and (14) can be expressed as*

$$\begin{aligned} f_k(x,t\,|\,c) &= \frac{1}{c+v}\sum_{j=0}^{k-2}\mathrm{P}\{N_{k-1} = j, Z_k = c\,|\,Z_0 = c\} \\ &\quad\times f_D^{(k-j-1)}(t-\tau_*)\int_{t-\tau_*}^{t} f_U^{(j+1)}(s-t+\tau_*)\overline{F}_{U_{j+2}}(t-s)\mathrm{d}s, \qquad k \ge 2, \end{aligned} \tag{15}$$

and

$$
\begin{aligned}
b_k(x,t\,|\,c) = \frac{1}{c+v}\Big\{&\mathrm{P}\{N_{k-1}=k-1, Z_k=-v\,|\,Z_0=c\} f_U^{(k)}(\tau_*)\overline{F}_{D_1}(t-\tau_*) \\
&+ \mathbf{1}_{\{k\geq 2\}}\sum_{j=0}^{k-2}\mathrm{P}\{N_{k-1}=j, Z_k=-v\,|\,Z_0=c\} \\
&\times f_U^{(j+1)}(\tau_*)\int_{\tau_*}^{t} f_D^{(k-j-1)}(s-\tau_*)\overline{F}_{D_{k-j}}(t-s)\mathrm{d}s\Big\}, \qquad k\geq 1,
\end{aligned} \tag{16}
$$

where

$$\tau_* = \tau_*(x,t) = \frac{vt+x}{c+v}, \tag{17}$$

and where $\mathrm{P}\{N_{k-1}=j, Z_k=c\,|\,Z_0=c\}$ *and* $\mathrm{P}\{N_{k-1}=j, Z_k=-v\,|\,Z_0=c\}$ *are given respectively by (45) and (46) if* $A=0$*, and by (50) and (51) if* $A>0$.

Proof. Recalling (13), for $t>0$, $-vt<x<ct$, and $k\geq 2$ we have

$$
\begin{aligned}
f_k(x,t\,|\,c)\mathrm{d}x = \int_0^t &\mathrm{P}\{T_k\in \mathrm{d}s, Z_k=c, S_s+c(t-s)\in \mathrm{d}x, T_{k+1}-T_k>t-s, \\
&0\leq N_{k-1}\leq k-2\,|\,Z_0=c\}.
\end{aligned}
$$

(Case $k=1$ does not give an absolutely continuous component). Conditioning on N_{k-1}, and taking into account the number of time periods during which the particle moved forward and backward, we obtain

$$
\begin{aligned}
f_k(x,t\,|\,c)\mathrm{d}x = \sum_{j=0}^{k-2}\int_0^t &\mathrm{P}\{U^{(j+1)}+D^{(k-j-1)}\in \mathrm{d}s, cU^{(j+1)}-vD^{(k-j-1)}+c(t-s)\in \mathrm{d}x\} \\
&\times \mathrm{P}\{U_{j+2}>t-s\}\mathrm{P}\{N_{k-1}=j, Z_k=c\,|\,Z_0=c\}.
\end{aligned}
$$

Note that conditions $S_s+c(t-s)=x$ and $S_s\geq -vs$ provide $s\geq (ct-x)/(c+v)=t-\tau_*$. This inequality and the independence of $U^{(k)}$ and $D^{(k)}$ thus give

$$f_k(x,t\,|\,c) = \sum_{j=0}^{k-2}\mathrm{P}\{N_{k-1}=j, Z_k=c\,|\,Z_0=c\}\int_{t-\tau_*}^{t} h(s,x-c(t-s))\mathrm{P}\{U_{j+2}>t-s\}\mathrm{d}s,$$

where $h(\cdot,\cdot)$ is the joint probability density of $(U^{(j+1)}+D^{(k-j-1)}, cU^{(j+1)}-vD^{(k-j-1)})$. Since

$$h(s,x-c(t-s)) = \frac{1}{c+v} f_U^{(j+1)}\left(s-\frac{ct-x}{c+v}\right) f_D^{(k-j-1)}\left(\frac{ct-x}{c+v}\right),$$

Eq. (15) thus follows recalling (17). Eq. (16) can be obtained in a similar way. Indeed for $k\geq 1$

$$
\begin{aligned}
b_k(x,t\,|\,c)\mathrm{d}x = \int_0^t &\mathrm{P}\{T_k\in \mathrm{d}s, Z_k=-v, S_s-v(t-s)\in \mathrm{d}x, T_{k+1}-T_k>t-s, \\
&0\leq N_{k-1}\leq k-1\,|\,Z_0=c\}.
\end{aligned}
$$

Conditioning on N_{k-1}, and taking into account the number of time periods during which the particle moved forward and backward, we obtain for $k \geq 1$

$$\begin{aligned}
b_k(x,t\,|\,c)\mathrm{d}x = {} & \int_0^t \mathrm{P}\{U^{(k)} \in \mathrm{d}s, cU^{(k)} - v(t-s) \in \mathrm{d}x\} \\
& \times \mathrm{P}\{D_1 > t-s\}\mathrm{P}\{N_{k-1} = k-1, Z_k = -v\,|\,Z_0 = c\} \\
& + \sum_{j=0}^{k-2} \int_0^t \mathrm{P}\{U^{(j+1)} + D^{(k-j-1)} \in \mathrm{d}s, cU^{(j+1)} - vD^{(k-j-1)} - v(t-s) \in \mathrm{d}x\} \\
& \times \mathrm{P}\{D_{k-j} > t-s\}\mathrm{P}\{N_{k-1} = j, Z_k = -v\,|\,Z_0 = c\}.
\end{aligned}$$

(Note that for $k = 1$ the above term $\mathrm{P}\{N_{k-1} = k-1, Z_k = -v\,|\,Z_0 = c\}$ reduces to $\mathrm{P}\{Z_1 = -v\,|\,Z_0 = c\}$ and the sum on j is equal to zero.) For the first integral, conditions $U^{(k)} = s$ and $cU^{(k)} - v(t-s) = x$ imply $s = \tau_*$; while in the second integral, conditions $S_s - v(t-s) = x$ and $S_s \leq cs$ provide $s \geq (x+vt)/(c+v) = \tau_*$. This inequality and the independence of $U^{(k)}$ and $D^{(k)}$ thus give for the above sum on j

$$\sum_{j=0}^{k-2} \mathrm{P}\{N_{k-1} = j, Z_k = -v\,|\,Z_0 = c\} \int_{\tau_*}^t h(s, x+v(t-s)) \mathrm{P}\{D_{k-j} > t-s\}\mathrm{d}s,$$

where, as above, $h(\cdot,\cdot)$ is the joint probability density of $(U^{(j+1)}+D^{(k-j-1)}, cU^{(j+1)}-vD^{(k-j-1)})$. Since

$$h(s, x+v(t-s)) = \frac{1}{c+v} f_U^{(j+1)}(\tau_*)\, f_D^{(k-j-1)}(s-\tau_*)\,,$$

we get (16). □

Remark 3.1 Similarly to (15) and (16), when the initial velocity is negative, for all $t > 0$ and $-vt < x < ct$, the densities (13) and (14) are expressed as

$$\begin{aligned}
f_k(x,t\,|-v) = {} & \frac{1}{c+v}\Big\{\mathrm{P}\{N_{k-1} = 0, Z_k = c\,|\,Z_0 = -v\} f_D^{(k)}(t-\tau_*)\overline{F}_{U_1}(\tau_*) \\
& + \mathbf{1}_{\{k\geq 2\}} \sum_{j=0}^{k-2} \mathrm{P}\{N_{k-1} = k-1-j, Z_k = c\,|\,Z_0 = -v\} \\
& \times f_D^{(j+1)}(t-\tau_*) \int_{t-\tau_*}^t f_U^{(k-j-1)}(s-t+\tau_*)\overline{F}_{U_{k-j}}(t-s)\mathrm{d}s\Big\}, \qquad k \geq 1,
\end{aligned}$$

$$\begin{aligned}
b_k(x,t\,|-v) = {} & \frac{1}{c+v} \sum_{j=0}^{k-2} \mathrm{P}\{N_{k-1} = k-1-j, Z_k = -v\,|\,Z_0 = -v\} \\
& \times f_U^{(k-j-1)}(\tau_*) \int_{\tau_*}^t f_D^{(j+1)}(s-\tau_*)\overline{F}_{D_{j+2}}(t-s)\mathrm{d}s, \qquad k \geq 2,
\end{aligned}$$

where τ_* is defined in (17) and where $\mathrm{P}\{N_{k-1} = k-1-j, Z_k = c\,|\,Z_0 = -v\}$ and $\mathrm{P}\{N_{k-1} = k-1-j, Z_k = -v\,|\,Z_0 = -v\}$ are given respectively by (47) and (48) if $A = 0$, and by (52) and (53) if $A > 0$.

In the following proposition, conditional on positive initial velocity, the mean of process V_t is expressed in terms of the cumulative distribution function of T_k. (Obviously, we can obtain a similar expression conditioning on negative initial velocity.) We remark that, by conditioning on the value of N_{k-1} and using the independence of $U^{(j+1)}$ and $D^{(k-j-1)}$, it follows:

$$F_{T_k|Z_0}(t\,|\,c) = \sum_{j=0}^{k-1} \mathrm{P}\{N_{k-1}=j\,|\,Z_0=c\}\mathrm{P}\{U^{(j+1)}+D^{(k-j-1)}\le t\}, \tag{18}$$

where

$$\mathrm{P}\{U^{(j+1)}+D^{(k-j-1)}\le t\} = \int_0^t F_D^{(k-j-1)}(t-s) f_U^{(j+1)}(s)\,\mathrm{d}s = \int_0^t F_U^{(j+1)}(t-s) f_D^{(k-j-1)}(s)\,\mathrm{d}s. \tag{19}$$

Proposition 3.2 *For all $t>0$ we have*

$$\mathrm{E}[V_t\,|\,V_0=c] = c\,\overline{F}_{U_1}(t) + c\pi_A \sum_{k=1}^{+\infty} \phi_k(t\,|\,c) + (-v)(1-\pi_A)\sum_{k=1}^{+\infty}\psi_k(t\,|\,c) \tag{20}$$

where $\pi_A = \dfrac{b+A}{b+A+r}$ *and*

$$\begin{aligned} \phi_k(t\,|\,c) &= F_{T_k|Z_0}(t\,|\,c) - \int_0^t F_{T_k|Z_0}(t-s\,|\,c) f_{U_{k+1}}(s)\mathrm{d}s, \\ \psi_k(t\,|\,c) &= F_{T_k|Z_0}(t\,|\,c) - \int_0^t F_{T_k|Z_0}(t-s\,|\,c) f_{D_{k+1}}(s)\mathrm{d}s. \end{aligned} \tag{21}$$

Proof. We recall that in both cases (by independence in the Bernoulli scheme and by exchangeability in Pólya scheme), we have for each $n\ge 2$

$$\mathrm{P}\{X_n=1\,|\,X_1=1\}=\pi_A, \qquad \mathrm{P}\{X_n=0\,|\,X_1=1\}=1-\pi_A\,. \tag{22}$$

Hence, for every positive integer $k\ge 1$, recalling (4) and the first of (5), we have

$$\mathrm{E}[V_t\,|\,V_0=c] = \mathrm{E}[Z_{M_t}\,|\,Z_0=c] = c\overline{F}_{U_1}(t) + \sum_{k=1}^{+\infty}\mathrm{E}[Z_k\cdot \mathbf{1}_{\{T_k\le t<T_{k+1}\}}\,|\,Z_0=c].$$

Since, for $k\ge 1$,

$$\begin{aligned} &\mathrm{E}[Z_k\cdot\mathbf{1}_{\{T_k\le t<T_{k+1}\}}\,|\,Z_0=c] \\ &\quad= c\,\mathrm{P}\{Z_k=c\,|\,Z_0=c\}\,\mathrm{E}[\mathbf{1}_{\{T_k\le t<T_{k+1}\}}\,|\,Z_0=c,Z_k=c] \\ &\quad+ (-v)\,\mathrm{P}\{Z_k=-v\,|\,Z_0=c\}\,\mathrm{E}[\mathbf{1}_{\{T_k\le t<T_{k+1}\}}\,|\,Z_0=c,Z_k=-v] \\ &\quad= c\,\mathrm{P}\{X_{k+1}=1\,|\,X_1=1\}\,\mathrm{P}\{T_k\le t<T_{k+1}\,|\,Z_0=c,Z_k=c\} \\ &\quad+ (-v)\mathrm{P}\{X_{k+1}=0\,|\,X_1=1\}\,\mathrm{P}\{T_k\le t<T_{k+1}\,|\,Z_0=c,Z_k=-v\} \\ &\quad= c\pi_A\int_0^\infty \mathrm{P}(t-s<T_k\le t\,|\,Z_0=c) f_{U_{k+1}}(s)\mathrm{d}s \\ &\quad+ (-v)(1-\pi_A)\int_0^\infty \mathrm{P}(t-s<T_k\le t\,|\,Z_0=c) f_{D_{k+1}}(s)\mathrm{d}s \\ &\quad= c\pi_A\,\phi_k(t\,|\,c) + (-v)(1-\pi_A)\,\psi_k(t\,|\,c)\,. \end{aligned}$$

Eq. (20) thus follows, due to (21). $\square$

The following remark turns out to be useful in some particular cases.

Remark 3.2 It is worthwhile to note that, using (18) and (19), Eqs. (21) can be rewritten as

$$\phi_k(t\,|\,c) = \sum_{j=0}^{k-1} \mathrm{P}\{N_{k-1}=j|Z_0=c\} \int_0^t F_D^{(k-j-1)}(t-y) \left[f_U^{(j+1)}(y) - f_{U^{(j+1)}+U_{k+1}}(y)\right] \mathrm{d}y,$$

$$\psi_k(t\,|\,c) = \sum_{j=0}^{k-1} \mathrm{P}\{N_{k-1}=j|Z_0=c\} \int_0^t F_U^{(j+1)}(t-y) \left[f_D^{(k-j-1)}(y) - f_{D^{(k-j-1)}+D_{k+1}}(y)\right] \mathrm{d}y.$$

For instance, the above quantities can be easily computed when the random variables U_k (and D_k) are gamma distributed with the same scale parameter, since it is well-known that, by independence, the sums of the involved random variables are still gamma distributed.

In the following sections we discuss some special cases arising in the two different schemes of Bernoulli and Pólya trials, and leading to closed forms for the probability law of S_t.

4 Particular case for Bernoulli scheme $(A = 0)$

The classical telegraph process is characterized by exponentially distributed times separating consecutive velocity changes. The extension of such a model to the case of velocities driven by Bernoulli trials can be performed in a simple and natural way. Indeed, if we consider a simple telegraph process $\tilde{S}_t$ with alternating velocities c and $-v$, and with alternating switching intensities $\tilde{\lambda} := (1-p)\lambda$ and $\tilde{\mu} := p\mu$, then it can be proved that the marginal distributions of $\tilde{S}_t$, $t \geq 0$, and S_t, $t \geq 0$, are identical (where S_t is a telegraph process with switching intensities λ and μ, and having the same velocities of $\tilde{S}_t$, governed by Bernoulli trials with parameter p). Hence, in this case results on S_t can be immediately obtained by those of $\tilde{S}_t$.

Therefore, aiming to discuss a non trivial case, and stimulated by previous studies (see Di Crescenzo and Martinucci [11] and Di Crescenzo *et al.* [12]) involving finite-velocity random motions with stochastically decreasing random intertimes, in the following we assume that the r.v.'s U_k and D_k have exponential distribution with linear rates λk and μk. Hence, the tail distribution functions are

$$\overline{F}_{U_k}(t) = \mathrm{e}^{-\lambda k t}, \qquad \overline{F}_{D_k}(t) = \mathrm{e}^{-\mu k t}, \qquad t \geq 0, \tag{23}$$

with $\lambda, \mu > 0$. Figure 2 shows some simulations of S_t in the present case, where the particle exhibits a kind of damped motion. Due to assumption (23), $U^{(k)}$ and $D^{(k)}$, $k \geq 1$, have generalized exponential densities

$$f_U^{(k)}(t) = k(1-\mathrm{e}^{-\lambda t})^{k-1}\lambda \mathrm{e}^{-\lambda t}, \qquad f_D^{(k)}(t) = k(1-\mathrm{e}^{-\mu t})^{k-1}\mu \mathrm{e}^{-\mu t}, \qquad t > 0 \tag{24}$$



Figure 2: Simulated sample paths of $\{S_t\}$ in the Bernoulli scheme, exponential damped case, with $\lambda = \mu = 1$ and $c = v = 1$, for some choices of p.

with corresponding cumulative distribution functions

$$F_U^{(k)}(t) = (1 - \mathrm{e}^{-\lambda t})^k, \qquad F_D^{(k)}(t) = (1 - \mathrm{e}^{-\mu t})^k, \qquad t \geq 0. \tag{25}$$

Hence, $U^{(k)}$ and $D^{(k)}$ are distributed as the maximum of k i.i.d. random variables having exponential distributions with rates λ and μ, respectively.

By making use of the results in the previous section, in the following we obtain the probability law of S_t. We start providing the discrete component.

Proposition 4.1 *Let U_k and D_k be exponentially distributed with rates λk and μk, $k = 1, 2, \cdots$, respectively. For all $t > 0$ we have*

$$\mathrm{P}\{S_t = ct\} = \frac{p\, e^{-\lambda t}}{1 - p\,(1 - e^{-\lambda t})}, \qquad \mathrm{P}\{S_t = -vt\} = \frac{(1-p)\, e^{-\mu t}}{1 - (1-p)\,(1 - e^{-\mu t})}.$$

Proof. It immediately follows from Proposition 3.1 and Eq. (25). ☐

The following theorem and corollary give the absolutely continuous component of the probability law of S_t.

Theorem 4.1 *Under the assumptions of Proposition 4.1, for all $t > 0$ and $-vt < x < ct$, we have*

$$f(x, t \,|\, c) = \frac{(1-p)p\mu \mathrm{e}^{\mu(t+\tau_*)}(\mathrm{e}^{\lambda\tau_*} - 1)}{(c+v)\left[p\mathrm{e}^{\mu t} + (1-p)\mathrm{e}^{(\lambda+\mu)\tau_*}\right]^2} \tag{26}$$

$$b(x, t \,|\, c) = \frac{(1-p)\left[\lambda p \mathrm{e}^{\lambda\tau_* + \mu(t+\tau_*)} + \lambda(1-p)\mathrm{e}^{\lambda\tau_* + 2\mu\tau_*}\right]}{(c+v)\left[p\mathrm{e}^{\mu t} + (1-p)\mathrm{e}^{(\lambda+\mu)\tau_*}\right]^2}, \tag{27}$$

where τ_* is defined in (17).

Proof. Since U_1 and U_{k+1} are exponentially distributed with parameters λ and $\lambda(k+1)$, respectively, from (12), (15) and (45), recalling (23) and (24) we get

$$\begin{aligned}
f(x,t\,|\,c) &= \frac{\mu}{c+v}\sum_{k=2}^{+\infty}\sum_{j=0}^{k-2}\binom{k-1}{j}p^{j+1}(1-p)^{k-1-j}(k-j-1)\mathrm{e}^{-\mu(t-\tau_*)}\left[1-\mathrm{e}^{-\mu(t-\tau_*)}\right]^{k-2-j} \\
&\quad\times\int_{t-\tau_*}^{t}\lambda(j+1)\mathrm{e}^{-\lambda(s-t+\tau_*)}\left[1-\mathrm{e}^{-\lambda(s-t+\tau_*)}\right]^{j}\mathrm{e}^{-\lambda(t-s)(j+2)}\mathrm{d}s \\
&= \frac{\mu p(1-p)}{c+v}\mathrm{e}^{-\mu(t-\tau_*)}\sum_{k=2}^{+\infty}(k-1)\sum_{j=0}^{k-2}\binom{k-2}{j}p^{j}(1-p)^{k-2-j}\left[1-\mathrm{e}^{-\mu(t-\tau_*)}\right]^{k-2-j} \\
&\quad\times\mathrm{e}^{-\lambda\tau_*}\int_{t-\tau_*}^{t}\lambda(j+1)\mathrm{e}^{-\lambda(t-s)}\left[\mathrm{e}^{-\lambda(t-s)}-\mathrm{e}^{-\lambda\tau_*}\right]^{j}\mathrm{d}s \\
&= \frac{\mu p(1-p)}{c+v}\mathrm{e}^{-\mu(t-\tau_*)-2\lambda\tau_*}(\mathrm{e}^{\lambda\tau_*}-1) \\
&\quad\times\sum_{k=2}^{+\infty}(k-1)\sum_{j=0}^{k-2}\binom{k-2}{j}\left[p(1-\mathrm{e}^{-\lambda\tau_*})\right]^{j}\left[(1-p)\left(1-\mathrm{e}^{-\mu(t-\tau_*)}\right)\right]^{k-j-2} \\
&= \frac{\mu p(1-p)}{c+v}\mathrm{e}^{-\mu(t-\tau_*)-2\lambda\tau_*}(\mathrm{e}^{\lambda\tau_*}-1)\,\frac{\mathrm{e}^{2\mu t+2\lambda\tau_*}}{\left[(1-p)\mathrm{e}^{(\lambda+\mu)\tau_*}+p\,\mathrm{e}^{\mu t}\right]^2}.
\end{aligned}$$

This yields density (26). Eq. (27) can be obtained from (12), (16) and (46) in a similar way. Indeed, for $k=1$ we have

$$b_1(x,t\,|\,c) = \frac{\lambda(1-p)}{c+v}\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}$$

and, for $k\geq 2$,

$$\begin{aligned}
b_k(x,t\,|\,c) &= \frac{1}{c+v}\left\{p^{k-1}(1-p)k\left(1-\mathrm{e}^{-\lambda\tau_*}\right)^{k-1}\lambda\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}+\right. \\
&\quad+\sum_{j=0}^{k-2}\binom{k-1}{j}p^{j}(1-p)^{k-j}(j+1)\left(1-\mathrm{e}^{-\lambda\tau_*}\right)^{j}\lambda\mathrm{e}^{-\lambda\tau_*} \\
&\quad\left.\times\int_{\tau_*}^{t}(k-j-1)\left(1-\mathrm{e}^{-\mu(s-\tau_*)}\right)^{k-j-2}\mu\mathrm{e}^{-\mu(s-\tau_*)-\mu(k-j)(t-s)}\,\mathrm{d}s\right\} \\
&= \frac{\lambda}{c+v}\left\{\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}(1-p)k\left[p(1-\mathrm{e}^{-\lambda\tau_*})\right]^{k-1}\right. \\
&\quad+\mathrm{e}^{-\lambda\tau_*}\sum_{j=0}^{k-2}\binom{k-1}{j}\left[p(1-\mathrm{e}^{-\lambda\tau_*})\right]^{j}(1-p)^{k-j}(j+1) \\
&\quad\left.\times\mathrm{e}^{-\mu(t-\tau_*)}\int_{\tau_*}^{t}(k-j-1)\left[\mathrm{e}^{-\mu(t-s)}-\mathrm{e}^{-\mu(t-\tau_*)}\right]^{k-j-2}\mu\mathrm{e}^{-\mu(t-s)}\,\mathrm{d}s\right\} \\
&= \frac{\lambda\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}(1-p)}{c+v}\left\{k\alpha^{k-1}+\sum_{j=0}^{k-2}\binom{k-1}{j}(j+1)\alpha^{j}\beta^{k-1-j}\right\},
\end{aligned}$$

where we have set

$$\alpha=\alpha(x,t)=p\left(1-\mathrm{e}^{-\lambda\tau_*}\right)\qquad\text{and}\qquad\beta=\beta(x,t)=(1-p)\left(1-\mathrm{e}^{-\mu(t-\tau_*)}\right).$$

Hence, recalling that (due to binomial theorem)

$$\sum_{j=0}^{k-2}\binom{k-1}{j}(j+1)\alpha^j\beta^{k-1-j}=(\alpha+\beta)^{k-1}+(k-1)\alpha(\alpha+\beta)^{k-2}-k\alpha^{k-1}\,,$$

from the second of (12) we finally get

$$\begin{aligned}b(x,t\,|\,c)&=\frac{\lambda\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}(1-p)}{c+v}\left\{1+\sum_{k=2}^{+\infty}(\alpha+\beta)^{k-1}+\alpha\sum_{k=2}^{+\infty}(k-1)(\alpha+\beta)^{k-2}\right\}\\&=\frac{\lambda\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}(1-p)}{c+v}\,\frac{1-\beta}{[1-(\alpha+\beta)]^2}\,,\end{aligned}$$

which coincides with Eq. (27). □

We are now able to find out the probability density of S_t in closed form.

Corollary 4.1 *Under the assumptions of Proposition 4.1, for all $t>0$ and $-vt<x<ct$, we have*

$$p(x,t\,|\,c)=\frac{1-p}{c+v}\,\frac{\lambda(1-p)\mathrm{e}^{(\lambda+2\mu)\tau_*}+p\,\mathrm{e}^{\mu(t+\tau_*)}\left[(\lambda+\mu)\mathrm{e}^{\lambda\tau_*}-\mu\right]}{\left[p\mathrm{e}^{\mu t}+(1-p)\mathrm{e}^{(\lambda+\mu)\tau_*}\right]^2}\tag{28}$$

and

$$p(x,t\,|-v)=\frac{p}{c+v}\,\frac{\mathrm{e}^{\mu(t+\tau_*)}\left[(\lambda+\mu)\mathrm{e}^{\lambda\tau_*}(1-p)+\mu p\right]-\lambda(1-p)\mathrm{e}^{(\lambda+2\mu)\tau_*}}{\left[p\mathrm{e}^{\mu t}+(1-p)\mathrm{e}^{(\lambda+\mu)\tau_*}\right]^2},\tag{29}$$

where τ_* is defined in (17). Hence, for $-vt<x<ct$ we have

$$p(x,t)=\frac{p}{1-p}\,\frac{1}{s}\,\frac{\exp\left\{(\mu-\frac{v}{s})t-\frac{1}{s}x\right\}}{\left[1+\frac{p}{1-p}\exp\left\{(\mu-\frac{v}{s})t-\frac{1}{s}x\right\}\right]^2},\tag{30}$$

where $s:=(c+v)/(\lambda+\mu)$.

Proof. Recalling (11), from densities (26) and (27) we obtain (28). By symmetry, density (29) can be expressed from (28) by replacing τ_* by $(t-\tau_*)$ and interchanging λ with μ and p with $(1-p)$. Therefore we finally get

$$p(x,t)=\frac{p(1-p)(\lambda+\mu)\,e^{\mu t+(\lambda+\mu)\tau_*}}{(c+v)[(1-p)\,e^{(\lambda+\mu)\tau_*}+pe^{\mu t}]^2},\qquad -vt<x<ct,$$

and then Eq. (30) easily follows. □

Some plots of density (30) are given in Figures 3, 4 and 5 for various choices of the involved parameters.

Remark 4.1 We can analyse the behavior of the density (30) when x tends to the endpoints of the state space $[-vt,ct]$. For $t>0$ we have

$$\lim_{x\downarrow -vt}p(x,t)=\frac{p}{1-p}\,\frac{1}{s}\,\frac{e^{\mu t}}{\left[1+\frac{p}{1-p}\,e^{\mu t}\right]^2},\qquad \lim_{x\uparrow ct}p(x,t)=\frac{p}{1-p}\,\frac{1}{s}\,\frac{e^{-\lambda t}}{\left[1+\frac{p}{1-p}\,e^{-\lambda t}\right]^2}.$$



Figure 3: Density (30), with $t = 1$, $c = v = 1$, $\lambda = 1$, for $\mu = 1$ (*left*) and $\mu = 2$ (*right*), and $p = 0.1,\ 0.2,\ 0.3,\ 0.4,\ 0.5$ (from bottom to top near $x = 1$).



Figure 4: Same as Figure 3, with $t = 10$ and $p = 0.1,\ 0.3,\ 0.5,\ 0.7,\ 0.9$ (from left to right in both plots).

The following special case follows by straightforward calculations.

Corollary 4.2 *Under the assumptions of Proposition 4.1, if $\lambda v = \mu c$ then density (30) becomes the truncated logistic density given by*

$$p(x,t) = \frac{e^{-(x-m)/s}}{s\left[1+e^{-(x-m)/s}\right]^2}, \qquad -vt < x < ct, \ \ t > 0,$$

$$m = s \ln\left(\frac{p}{1-p}\right), \qquad s = \frac{v}{\mu} = \frac{c}{\lambda}.$$

Hence, in this case S_t admits of the stationary density

$$\lim_{t\to+\infty} p(x,t) = \frac{e^{-(x-m)/s}}{s\left[1+e^{-(x-m)/s}\right]^2}, \qquad x \in \mathbb{R}. \tag{31}$$

We note that the right-hand-side of Eq. (31) is a logistic density with mean m and variance $\pi^2 s^2/3$. In addition, if $p = 1/2$ then the mean m vanishes, and the density identifies with the stationary p.d.f. of a damped telegraph process, as obtained in Corollary 3.3 of Di Crescenzo



Figure 5: Same as Figure 4, for $p = 0.1$ (*left*) and $p = 0.5$ (*right*), with $\mu = 1$, 2, 3, 4 (from left to right in both plots).

and Martinucci [11]. We remark that if $\lambda\, v \neq \mu\, c$ then

$$\lim_{t\to+\infty} p(x,t) = 0, \qquad x \in \mathbb{R}.$$

Finally, let us obtain the mean velocity conditioned by positive initial velocity.

Proposition 4.2 *Under the assumptions of Proposition 4.1, for all $t > 0$ we have*

$$\mathrm{E}[V_t \,|\, V_0 = c] = c\,\mathrm{e}^{-\lambda t} + c\pi_A \sum_{k=1}^{+\infty} \phi_k(t\,|\,c) + (-v)(1-\pi_A) \sum_{k=1}^{+\infty} \psi_k(t\,|\,c),$$

where

$$\begin{aligned}\phi_k(t\,|\,c) &= \lambda \sum_{j=0}^{k-1} (j+1)\binom{k-1}{j} p^j (1-p)^{k-1-j} \sum_{\ell=0}^{k-j-1} \sum_{h=0}^{j} \binom{k-j-1}{\ell}\binom{j}{h}(-1)^{\ell+h} \\ &\quad\times \Big[t\mathrm{e}^{-\mu\ell t}\, {}_1F_1(1,2;[\mu\ell-\lambda(h+1)]t) \\ &\quad -\lambda(k+1)\,\mathrm{e}^{-\lambda(h+1)t}\, H(\mu\ell-\lambda(h+1),\lambda(k-h);t)\Big], \\ \psi_k(t\,|\,c) &= \lambda \sum_{j=0}^{k-1} (j+1)\binom{k-1}{j} p^j (1-p)^{k-1-j} \sum_{\ell=0}^{k-j-1} \sum_{h=0}^{j} \binom{k-j-1}{\ell}\binom{j}{h}(-1)^{\ell+h} \\ &\quad\times \Big[t\mathrm{e}^{-\mu\ell t}\, {}_1F_1(1,2;[\mu\ell-\lambda(h+1)]t) \\ &\quad -\mu(k+1)\,\mathrm{e}^{-\lambda(h+1)t}\, H(\mu\ell-\lambda(h+1),\mu(k+1)-\lambda(h+1);t)\Big],\end{aligned}$$

and where the function H is defined by (59) in Appendix B.

Proof. The above formulas easily follow from Theorem 3.2. Indeed, after some calculations, from (18) and (19) we find the following cumulative distribution function:

$$\begin{aligned}F_{T_k|Z_0}(t\,|\,c) &= \lambda \sum_{j=0}^{k-1} (j+1)\binom{k-1}{j} p^j (1-p)^{k-1-j} \sum_{\ell=0}^{k-j-1} \sum_{h=0}^{j} \binom{k-j-1}{\ell}\binom{j}{h}(-1)^{\ell+h} \\ &\quad\times t\mathrm{e}^{-\mu\ell t}\, {}_1F_1(1,2;[\mu\ell-\lambda(h+1)]t),\end{aligned}$$

where ${}_1F_1(1,2;0)=1$ and ${}_1F_1(1,2;z)=(\mathrm{e}^z-1)/z$, for $z\neq 0$. Moreover, the thesis follows by computing the two integrals which appear in the right-hand-sides of Eqs. (21). $\square$

5 Particular case for Pólya scheme $(A>0)$

In this section we consider a special case in which the velocity changes are governed by the Pólya urn scheme. Let us assume the distributions of U_1 and D_1 are $\Gamma\left(\frac{b}{A}+1,\lambda\right)$ and $\Gamma\left(\frac{r}{A}+1,\mu\right)$, respectively, and the intertimes U_k and D_k, for $k\geq 2$, are exponential with parameters λ and μ, respectively. Therefore, $U^{(k)}$ has Gamma distribution $\Gamma\left(\frac{b}{A}+k,\lambda\right)$ and $D^{(k)}$ has Gamma distribution $\Gamma\left(\frac{r}{A}+k,\mu\right)$, so that

$$f_U^{(k)}(t)=\frac{\lambda^{b/A+k}\,t^{b/A+k-1}\,\mathrm{e}^{-\lambda t}}{\Gamma\left(\frac{b}{A}+k\right)},\qquad f_D^{(k)}(t)=\frac{\mu^{r/A+k}\,t^{r/A+k-1}\,\mathrm{e}^{-\mu t}}{\Gamma\left(\frac{r}{A}+k\right)},\qquad t>0. \tag{32}$$

Notice that U_1 (resp., D_1) is stochastically larger than U_k (resp., D_k), $k\geq 2$. Hence, the first time interval along both directions is stochastically greater than the other ones in the same direction. This is not an unusual assumption, since in renewal theory the distribution of the first interarrival time is often supposed different from that of the other ones (see Chapter 2 of Wolff [25], for instance), as for the delayed renewal processes. Moreover, differently from the case treated in the previous section, U_1 (resp., D_1) depends on b and A (resp., r and A), which are the same parameters involved in the Pólya urn scheme described in Section 2.

Under the above assumptions we are able to explicitly obtain the probability law of S_t in terms of the hypergeometric function ${}_1F_1(u,v;z)$ (see Eq. (54) in Appendix B). We first provide the tail distribution functions of $U^{(k)}$ and $D^{(k)}$ (see Eq. (57) in Appendix B):

$$\overline{F}_U^{(k)}(t)=1-\frac{(\lambda t)^{b/A+k}\mathrm{e}^{-\lambda t}}{\Gamma\left(\frac{b}{A}+k+1\right)}{}_1F_1\left(1,\frac{b}{A}+k+1;\lambda t\right),\qquad t\geq 0, \tag{33}$$

$$\overline{F}_D^{(k)}(t)=1-\frac{(\mu t)^{r/A+k}\mathrm{e}^{-\mu t}}{\Gamma\left(\frac{r}{A}+k+1\right)}{}_1F_1\left(1,\frac{r}{A}+k+1;\mu t\right),\qquad t\geq 0\,. \tag{34}$$

In the following proposition we give the discrete component of the probability law of S_t.

Proposition 5.1 *Let U_1 and D_1 be gamma distributed with parameters $\left(\frac{b}{A}+1,\lambda\right)$ and $\left(\frac{r}{A}+1,\mu\right)$, respectively, and let U_k and D_k, for $k\geq 2$, be exponentially distributed with parameters λ and μ, respectively. For all $t>0$ we have*

$$\mathrm{P}\{S_t=ct\}=\frac{b}{b+r}\overline{F}_{U_1}(t)+\frac{b\,(\lambda t)^{b/A}\,\mathrm{e}^{-\lambda t}}{(b+r)\,\Gamma\left(\frac{b+A}{A}\right)}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda t\right)-1\right], \tag{35}$$

where $\overline{F}_{U_1}(t)$ is given by (33) for $k=1$. Similarly,

$$\mathrm{P}\{S_t=-vt\}=\frac{r}{b+r}\overline{F}_{D_1}(t)+\frac{r\,(\mu t)^{r/A}\,\mathrm{e}^{-\mu t}}{(b+r)\,\Gamma\left(\frac{r+A}{A}\right)}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\mu t\right)-1\right], \tag{36}$$

where $\overline{F}_{D_1}(t)$ is given by (34) for $k=1$.

Proof. Eq. (35) follows from (7) using (32). Indeed, since

$$F_U^{(k)}(t) - F_U^{(k+1)}(t) = \int_0^t f_U^{(k)}(s) \overline{F}_{U_{k+1}}(t-s)\,\mathrm{d}s\,,$$

we obtain

$$\begin{aligned}
\mathrm{P}\{S_t = ct\} &= \frac{b}{b+r}\overline{F}_{U_1}(t) + \sum_{k=1}^{+\infty} \mathrm{P}\{Z_0 = c, N_k = k\} \int_0^t \lambda^{b/A+k} \frac{s^{b/A+k-1}\mathrm{e}^{-\lambda s}}{\Gamma\left(\frac{b}{A}+k\right)} \mathrm{e}^{-\lambda(t-s)} \mathrm{d}s \\
&= \frac{b}{b+r}\overline{F}_{U_1}(t) + \frac{b\mathrm{e}^{-\lambda t}\,(\lambda t)^{b/A}}{(b+r)\Gamma\left(\frac{b+A}{A}\right)} \sum_{k=1}^{+\infty} \frac{(1)_k}{\left(\frac{b+A+r}{A}\right)_k} \frac{(\lambda t)^k}{k!} \\
&= \frac{b}{b+r}\overline{F}_{U_1}(t) + \frac{b\mathrm{e}^{-\lambda t}\,(\lambda t)^{b/A}}{(b+r)\Gamma\left(\frac{b+A}{A}\right)} \left[{}_1F_1\left(1, \frac{b+A+r}{A}; \lambda t\right) - 1\right].
\end{aligned}$$

By interchanging λ with μ, b with r, and U with D in (35), we immediately obtain (36). $\square$

For the absolutely continuous component of the probability law of S_t we have the following results.

Theorem 5.1 *Under the assumptions of Proposition 5.1, for all $t > 0$ and $-vt < x < ct$, we have*

$$f(x,t\,|\,c) = \frac{\xi(\tau_*,t)\eta(\tau_*,t)}{t-\tau_*} \tag{37}$$

and

$$b(x,t\,|\,c) = \frac{r\lambda(\lambda\tau_*)^{b/A-1}\mathrm{e}^{-\lambda\tau_*}}{(c+v)A\Gamma\left(\frac{b+A}{A}\right)} \overline{F}_{D_1}(t-\tau_*) \left[{}_1F_1\left(1, \frac{b+A+r}{A}; \lambda\tau_*\right) - 1\right] + \frac{\xi(\tau_*,t)\eta(\tau_*,t)}{\tau_*}, \tag{38}$$

where τ_* is defined in (17), and where

$$\xi(\tau_*,t) := \exp\{-\lambda\tau_* - \mu(t-\tau_*)\} \frac{(\lambda\tau_*)^{b/A+1}\left[\mu(t-\tau_*)\right]^{r/A}}{(c+v)\,\Gamma\left(\frac{b+A}{A}\right)\Gamma\left(\frac{r}{A}\right)}\,, \tag{39}$$

$$\begin{aligned}
\eta(\tau_*,t) := {} & [\lambda\tau_* + \mu(t-\tau_*)]^{-1} \left[{}_1F_1\left(1, \frac{b+A+r}{A}; \lambda\tau_* + \mu(t-\tau_*)\right) - 1 - \frac{A\left[\lambda\tau_* + \mu(t-\tau_*)\right]}{b+A+r}\right] \\
& - (\lambda\tau_*)^{-1} \left[{}_1F_1\left(1, \frac{b+A+r}{A}; \lambda\tau_*\right) - 1 - \frac{A\lambda\tau_*}{b+A+r}\right].
\end{aligned} \tag{40}$$

Proof. Due to Eqs. (12), (16) and (50), and recalling (32), under the given assumptions we have

$$
\begin{aligned}
&f(x,t\,|\,c)=\frac{1}{c+v}\sum_{k=2}^{+\infty}\sum_{j=0}^{k-2}\mathrm{P}\{N_{k-1}=j,Z_k=c\,|\,Z_0=c\}\\
&\times\int_{t-\tau_*}^{t}\frac{\lambda^{b/A+j+1}(s-t+\tau_*)^{b/A+j}}{\Gamma\left(\frac{b}{A}+j+1\right)}\mathrm{e}^{-\lambda(s-t+\tau_*)}\frac{\mu^{r/A+k-j-1}(t-\tau_*)^{r/A+k-j-2}}{\Gamma\left(\frac{r}{A}+k-j-1\right)}\mathrm{e}^{-\mu(t-\tau_*)}\mathrm{e}^{-\lambda(t-s)}\mathrm{d}s\\
&=\frac{\xi(\tau_*,t)}{t-\tau_*}\sum_{k=2}^{+\infty}\frac{1}{\left(\frac{b+A+r}{A}\right)_k}\sum_{j=0}^{k-2}\binom{k-1}{j}(\lambda\tau_*)^j[\mu(t-\tau_*)]^{k-1-j}\\
&=\frac{\xi(\tau_*,t)}{t-\tau_*}\sum_{k=2}^{+\infty}\frac{(1)_k}{\left(\frac{b+A+r}{A}\right)_k}\frac{1}{k!}\left\{[\lambda\tau_*+\mu(t-\tau_*)]^{k-1}-(\lambda\tau_*)^{k-1}\right\}\\
&=\frac{\xi(\tau_*,t)}{t-\tau_*}\left\{[\lambda\tau_*+\mu(t-\tau_*)]^{-1}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*+\mu(t-\tau_*)\right)-1-\frac{A\left[\lambda\tau_*+\mu(t-\tau_*)\right]}{b+A+r}\right]\right.\\
&\left.-(\lambda\tau_*)^{-1}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*\right)-1-\frac{A\lambda\tau_*}{b+A+r}\right]\right\}.
\end{aligned}
$$

Similarly, Eqs. (12), (16) and (51) give

$$
\begin{aligned}
b(x,t\,|\,c)=\frac{1}{c+v}&\left\{\sum_{k=1}^{+\infty}\mathrm{P}\{N_{k-1}=k-1,Z_k=-v\,|\,Z_0=c\}\frac{\lambda^{b/A+k}\tau_*^{b/A+k-1}\mathrm{e}^{-\lambda\tau_*}}{\Gamma\left(\frac{b}{A}+k\right)}\overline{F}_{D_1}(t-\tau_*)\right.\\
&+\sum_{k=2}^{+\infty}\sum_{j=0}^{k-2}\mathrm{P}\{N_{k-1}=j,Z_k=-v\,|\,Z_0=c\}\\
&\left.\times\frac{\lambda^{b/A+j+1}\tau_*^{b/A+j}\mu^{r/A+k-j-1}\mathrm{e}^{-\lambda\tau_*-\mu(t-\tau_*)}}{\Gamma\left(\frac{b}{A}+j+1\right)\Gamma\left(\frac{r}{A}+k-j-1\right)}\int_{\tau_*}^{t}(s-\tau_*)^{r/A+k-j-2}\mathrm{d}s\right\}\\
&=\frac{r(\lambda\tau_*)^{b/A}}{A\Gamma\left(\frac{b+A}{A}\right)\tau_*(c+v)}\mathrm{e}^{-\lambda\tau_*}\overline{F}_{D_1}(t-\tau_*)\sum_{k=1}^{+\infty}\frac{(1)_k}{\left(\frac{b+A+r}{A}\right)_k}\frac{(\lambda\tau_*)^k}{k!}\\
&+\frac{\xi(\tau_*,t)}{\tau_*}\sum_{k=2}^{+\infty}\frac{1}{\left(\frac{b+A+r}{A}\right)_k}\sum_{j=0}^{k-2}\binom{k-1}{j}(\lambda\tau_*)^j[\mu(t-\tau_*)]^{k-1-j}\\
&=\frac{r(\lambda\tau_*)^{b/A}}{A\Gamma\left(\frac{b+A}{A}\right)\tau_*(c+v)}\mathrm{e}^{-\lambda\tau_*}\overline{F}_{D_1}(t-\tau_*)\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*\right)-1\right]\\
&+\frac{\xi(\tau_*,t)}{\tau_*}\sum_{k=2}^{+\infty}\frac{(1)_k}{\left(\frac{b+A+r}{A}\right)_k}\frac{1}{k!}\left[[\lambda\tau_*+\mu(t-\tau_*)]^{k-1}-(\lambda\tau_*)^{k-1}\right]\\
&=\frac{r(\lambda\tau_*)^{b/A}}{A\Gamma\left(\frac{b+A}{A}\right)\tau_*(c+v)}\mathrm{e}^{-\lambda\tau_*}\overline{F}_{D_1}(t-\tau_*)\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*\right)-1\right]+\frac{\xi(\tau_*,t)}{\tau_*}\\
&\times\left\{[\lambda\tau_*+\mu(t-\tau_*)]^{-1}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*+\mu(t-\tau_*)\right)-1-\frac{A[\lambda\tau_*+\mu(t-\tau_*)]}{b+A+r}\right]\right.\\
&\left.-(\lambda\tau_*)^{-1}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*\right)-1-\frac{A(\lambda\tau_*)}{b+A+r}\right]\right\}.
\end{aligned}
$$

The proof is thus complete. □

Corollary 5.1 *Under the assumptions of Theorem 5.1, for all* $t > 0$ *and* $-vt < x < ct$*, the probability density of* S_t *is given by Eq. (10), where*

$$p(x,t\,|\,c) = \frac{r\lambda(\lambda\tau_*)^{b/A-1}\mathrm{e}^{-\lambda\tau_*}}{(c+v)A\Gamma\left(\frac{b+A}{A}\right)}\overline{F}_{D_1}(t-\tau_*)\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*\right)-1\right]+\frac{\xi(\tau_*,t)\eta(\tau_*,t)\,t}{\tau_*(t-\tau_*)}\,, \tag{41}$$

$$\begin{aligned}p(x,t\,|-v) = {}& \frac{b\mu\big[\mu(t-\tau_*)\big]^{r/A-1}\mathrm{e}^{-\mu(t-\tau_*)}}{(c+v)A\Gamma\left(\frac{r+A}{A}\right)}\overline{F}_{U_1}(\tau_*)\left[{}_1F_1\left(1,\frac{b+A+r}{A};\mu(t-\tau_*)\right)-1\right]\\ &+\frac{\widetilde{\xi}(\tau_*,t)\widetilde{\eta}(\tau_*,t)\,t}{\tau_*(t-\tau_*)}\,,\end{aligned} \tag{42}$$

where τ_*, ξ *and* η*, are defined respectively in Eqs. (17), (39) and (40), and where*

$$\widetilde{\xi}(\tau_*,t) := \exp\{-\lambda\tau_* - \mu(t-\tau_*)\}\,\frac{(\lambda\tau_*)^{b/A}\big[\mu(t-\tau_*)\big]^{r/A+1}}{(c+v)\Gamma\left(\frac{r+A}{A}\right)\Gamma\left(\frac{b}{A}\right)}\,,$$

$$\begin{aligned}\widetilde{\eta}(\tau_*,t) := {}& [\lambda\tau_*+\mu(t-\tau_*)]^{-1}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\lambda\tau_*+\mu(t-\tau_*)\right)-1-\frac{A\,[\lambda\tau_*+\mu(t-\tau_*)]}{b+A+r}\right]\\ &-[\mu(t-\tau_*)]^{-1}\left[{}_1F_1\left(1,\frac{b+A+r}{A};\mu(t-\tau_*)\right)-1-\frac{A\mu(t-\tau_*)}{b+A+r}\right].\end{aligned}$$

Proof. Eq. (41) immediately follows from (11). From Remark 3.1, by interchanging μ with λ, $(t-\tau_*)$ with τ_*, b with r, and D with U in $b(x,t\,|\,c)$, given in (38), we get the following density:

$$\begin{aligned}f(x,t\,|-v) = {}& \frac{b\mu\big[\mu(t-\tau_*)\big]^{r/A-1}}{(c+v)A\Gamma\left(\frac{r+A}{A}\right)}\mathrm{e}^{-\mu(t-\tau_*)}\overline{F}_{U_1}(\tau_*)\left[{}_1F_1\left(1,\frac{b+A+r}{A};\mu(t-\tau_*)\right)-1\right]\\ &+\frac{\widetilde{\xi}(\tau_*,t)\widetilde{\eta}(\tau_*,t)}{t-\tau_*}.\end{aligned}$$

Similarly, we obtain the density $b(x,t\,|-v)$ by interchanging μ with λ, $(t-\tau_*)$ with τ_*, b with r and D with U in $f(x,t\,|\,c)$, given in (37), so that

$$b(x,t\,|-v) = \frac{\widetilde{\xi}(\tau_*,t)\widetilde{\eta}(\tau_*,t)}{\tau_*}\,.$$

Hence, Eq. (42) follows by (11). □

Various plots of density $p(x,t)$ given in Corollary 5.1 are shown by Figures 6, 7 and 8.

Remark 5.1 From the formulas given in Theorem 5.1 we have

$$\lim_{\tau_*\to 0}\frac{\xi(\tau_*,t)\eta(\tau_*,t)}{t-\tau_*} = 0, \qquad \lim_{\tau_*\to t}\frac{\xi(\tau_*,t)\eta(\tau_*,t)}{t-\tau_*} = 0,$$

$$\lim_{\tau_*\to 0}\frac{\xi(\tau_*,t)\eta(\tau_*,t)}{\tau_*} = 0, \qquad \lim_{\tau_*\to t}\frac{\xi(\tau_*,t)\eta(\tau_*,t)}{\tau_*} = 0.$$



Figure 6: Density $p(x,t)$ for Pólya scheme with densities (32), with $t = 1$, $c = v = 1$, $\lambda = \mu = 1$, $A = 2$, for $b = 1$ (*left*) and $b = 2$ (*right*), and $r = 1, 2, 3, 4$ (from bottom to top near $x = 0$).



Figure 7: Same as Figure 6, with $r = 1$ and $A = 0.4, 0.6, 0.8, 1$ (from bottom to top near $x = 0$).

Hence, some calculations allow to study the behaviour of the density $p(x,t)$ in proximity of the endpoints $-vt$, ct:

$$\lim_{x\uparrow -vt} p(x,t) = \frac{br\mu(\mu t)^{r/A-1}\mathrm{e}^{-\mu t}}{(b+r)(c+v)A\Gamma(r/A+1)}\left[{}_1F_1\left(1, \frac{b+A+r}{A}; \mu t\right) - 1\right]$$

and

$$\lim_{x\uparrow ct} p(x,t) = \frac{br\lambda(\lambda t)^{b/A-1}\mathrm{e}^{-\lambda t}}{(b+r)(c+v)A\Gamma(b/A+1)}\left[{}_1F_1\left(1, \frac{b+A+r}{A}; \lambda t\right) - 1\right].$$

Note that in this case a stationary distribution does not exist. Indeed,

$$\lim_{t\to+\infty} \mathrm{P}\{S_t = ct\} = 0, \qquad \lim_{t\to+\infty} \mathrm{P}\{S_t = -vt\} = 0, \qquad \lim_{t\to+\infty} p(x,t) = 0,$$

since (see Equation 13.1.4 of [1]), as $|z| \to +\infty$,

$${}_1F_1(a,b;z) = \frac{\Gamma(b)}{\Gamma(a)}\,\mathrm{e}^z z^{a-b}[1 + O(|z|^{-1})] \qquad \text{for } \mathrm{Re}(z) > 0\,.$$

Finally, the mean velocity subordinated to the positive initial velocity can be expressed in the following form.



Figure 8: Same as Figure 6 (from bottom to top near $x = -8$), with $t = 10$.

Proposition 5.2 *Under the assumptions of Theorem 5.1, for all* $t > 0$ *we have*

$$\mathrm{E}[V_t \,|\, V_0 = c] = c\,\overline{F}_{U_1}(t) + c\pi_A \sum_{k=1}^{+\infty} \phi_k(t \,|\, c) + (-v)(1-\pi_A) \sum_{k=1}^{+\infty} \psi_k(t \,|\, c) \tag{43}$$

with

$$\begin{aligned} \phi_k(t \,|\, c) &= \sum_{j=0}^{k-1} \mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\} \\ &\quad \times \left[G(r/A+k-j-1, \mu, b/A+j+1, \lambda; t) - G(r/A+k-j-1, \mu, b/A+j+2, \lambda; t)\right] , \\ \psi_k(t \,|\, c) &= \sum_{j=0}^{k-1} \mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\} \\ &\quad \times \left[G(b/A+j+1, \lambda, r/A+k-j-1, \mu; t) - G(b/A+j+1, \lambda, r/A+k-j, \mu; t)\right] , \end{aligned} \tag{44}$$

where $\mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\}$ *is given by (49) and the function* G *is defined in Appendix B, by Eq. (58).*

Proof. Eqs. (43) and (44) easily follow from Proposition 3.2, Remark 3.2 and Eq. (58) since, for $k \geq 1$, U_{k+1} and D_{k+1} are exponential with parameter λ and μ, respectively, and $U^{(k)}$ and $D^{(k)}$ are Gamma-distributed with parameters $(b/A + k, \lambda)$ and $(r/A + k, \mu)$ respectively. ☐

5.1 A suitable extension

In this section we propose a suitable extension of the model based on Pólya scheme by replacing the classical Pólya urn with the following randomly reinforced urn (cf. Aletti *et al.* [2], Berti *et al.* [5], [6], Crimaldi [7]). An urn contains $b > 0$ black balls and $r > 0$ red balls. At each epoch $n \geq 1$, a ball is drawn and then replaced together with a random number of balls of the same color. Say that $B_n \geq 0$ black balls or $R_n \geq 0$ red balls are added to the urn according to whether $X_n = 1$ or $X_n = 0$, where X_n is the indicator function of the event {black ball at time n}. We assume, for each $n \geq 1$,

$$(B_n, R_n) \text{ independent of } \big(X_1, B_1, R_1, X_2, \ldots, B_{n-1}, R_{n-1}, X_n\big).$$

According to this new model, formula (2) must be replaced by

$$\begin{aligned} \mathrm{P}\{Z_n = c \,|\, \mathcal{G}_n\} = \mathrm{P}\{X_{n+1} = 1 \,|\, \mathcal{G}_n\} &= \frac{b + \sum_{k=1}^n B_k X_k}{b + r + \sum_{k=1}^n \big(B_k X_k + R_k(1 - X_k)\big)}, \\ \mathrm{P}\{Z_n = -v \,|\, \mathcal{G}_n\} = \mathrm{P}\{X_{n+1} = 0 \,|\, \mathcal{G}_n\} &= \frac{r + \sum_{k=1}^n R_k(1 - X_k)}{b + r + \sum_{k=1}^n \big(B_k X_k + R_k(1 - X_k)\big)}, \end{aligned}$$

where

$$\mathcal{G}_0 = \{\emptyset, \Omega\}, \qquad \mathcal{G}_n = \sigma\big(X_1, B_1, R_1, \ldots, X_n, B_n, R_n\big).$$

It has been proved that, if $B_n = R_n$ for each n (cf. Berti et al. [6] and Crimaldi [7]) or, more generally, if[2] $B_n \overset{d}{=} R_n$ for each n (cf. Berti et al. [5], p. 538), then $(X_n)_{n \geq 1}$ is a sequence of conditionally identically distributed random variables (which is generally not exchangeable) and so we have, for $n \geq 2$,

$$\mathrm{P}\{X_n = 1 \,|\, X_1 = 1\} = \mathrm{E}\left[\frac{b + B_1}{b + B_1 + r}\right], \qquad \mathrm{P}\{X_n = 0 \,|\, X_1 = 1\} = \mathrm{E}\left[\frac{r}{b + B_1 + r}\right].$$

Using the above equalities, instead of (22), we can obtain a formula for the mean velocity similar to (20). However, in this case the computation of the conditional probability distribution of N_{k-1} needed in Eq. (18) is not an easy task, and possibly will be the object of future investigations.

A The conditional probability distribution of N_{k-1}

A.1 Bernoulli scheme ($A = 0$)

Using the independence of $\{X_n; n \geq 1\}$ and setting p as in (3), we have for $k \geq 1$ and $0 \leq j \leq k - 1$

$$\mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\} = \mathrm{P}\left\{\sum_{h=2}^{k} X_h = j \,|\, X_1 = 1\right\} = \binom{k-1}{j} p^j (1 - p)^{k-1-j}$$

and so

$$\mathrm{P}\{N_{k-1} = j,\, Z_k = c \,|\, Z_0 = c\} = \binom{k-1}{j} p^{j+1} (1 - p)^{k-1-j}\,, \tag{45}$$

which simply becomes $\mathrm{P}\{Z_1 = c \,|\, Z_0 = c\} = p$ for $k = 1$ and $j = 0$. Similarly, we have

$$\mathrm{P}\{N_{k-1} = j,\, Z_k = -v \,|\, Z_0 = c\} = \binom{k-1}{j} p^j (1 - p)^{k-j}\,. \tag{46}$$

---

[2] The notation $\overset{d}{=}$ denotes equality in distribution

Conditioning on $Z_0 = -v$, we get

$$\mathrm{P}\{N_{k-1} = k-1-j,\, Z_k = c \,|\, Z_0 = -v\} = \binom{k-1}{j} p^{k-j}(1-p)^j \tag{47}$$

and

$$\mathrm{P}\{N_{k-1} = k-1-j,\, Z_k = -v \,|\, Z_0 = -v\} = \binom{k-1}{j} p^{k-1-j}(1-p)^{j+1}\,. \tag{48}$$

A.2 Pólya scheme $(A > 0)$

Using the exchangeability of $\{X_n; n \geq 1\}$, we have for $k \geq 1$ and $0 \leq j \leq k-1$

$$\begin{aligned}\mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\} &= \binom{k-1}{j} \frac{\Gamma\left(\frac{b+A+r}{A}\right)}{\Gamma\left(\frac{b+A}{A}\right)\Gamma\left(\frac{r}{A}\right)} \frac{\Gamma\left(j + \frac{b+A}{A}\right)\Gamma\left(k-1-j+\frac{r}{A}\right)}{\Gamma\left(k-1+\frac{b+A+r}{A}\right)}\,,\\ &= \binom{k-1}{j}\left(\frac{b+A}{A}\right)_j \left(\frac{r}{A}\right)_{k-j-1}\left[\left(\frac{b+A+r}{A}\right)_{k-1}\right]^{-1},\end{aligned} \tag{49}$$

where $(\alpha)_0 = 1$ and $(\alpha)_j = \alpha(\alpha+1)\cdot\ldots\cdot(\alpha+j-1) = \Gamma(\alpha+j)/\Gamma(\alpha)$ (the ascending factorial). For $k \geq 2$, the above formula corresponds to the quantity $\mathrm{P}\{\sum_{h=2}^{k} X_h = j \,|\, X_1 = 1\}$, i.e. to the probability of obtaining j black balls in $k-1$ drawings in a Pólya urn scheme starting from r red balls and $b + A$ black balls. Hence we get

$$\begin{aligned}\mathrm{P}\{N_{k-1} = j,\, Z_k = c \,|\, Z_0 = c\} &= \mathrm{P}\{X_{k+1} = 1 \,|\, N_{k-1} = j,\, X_1 = 1\} \cdot \mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\}\\ &= \frac{b + A(j+1)}{b+r+Ak}\mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\}\\ &= \binom{k-1}{j}\frac{\Gamma\left(\frac{b+A+r}{A}\right)\Gamma\left(j+1+\frac{b+A}{A}\right)\Gamma\left(k-1-j+\frac{r}{A}\right)}{\Gamma\left(\frac{b+A}{A}\right)\Gamma\left(\frac{r}{A}\right)\Gamma\left(k+\frac{b+A+r}{A}\right)}\\ &= \binom{k-1}{j}\left(\frac{b+A}{A}\right)_{j+1}\left(\frac{r}{A}\right)_{k-j-1}\left[\left(\frac{b+A+r}{A}\right)_k\right]^{-1},\end{aligned} \tag{50}$$

which simply becomes $\mathrm{P}\{Z_1 = c \,|\, Z_0 = c\} = \frac{b+A}{b+r+A}$ for $k = 1$ and $j = 0$. Similarly, we have

$$\begin{aligned}\mathrm{P}\{N_{k-1} = j,\, Z_k = -v \,|\, Z_0 = c\} &= \mathrm{P}\{X_{k+1} = 0 \,|\, N_{k-1} = j,\, X_1 = 1\} \cdot \mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\}\\ &= \frac{r + A(k-j-1)}{b+r+Ak}\mathrm{P}\{N_{k-1} = j \,|\, Z_0 = c\}\\ &= \binom{k-1}{j}\frac{\Gamma\left(\frac{b+A+r}{A}\right)\Gamma\left(j+\frac{b+A}{A}\right)\Gamma\left(k-j+\frac{r}{A}\right)}{\Gamma\left(\frac{b+A}{A}\right)\Gamma\left(\frac{r}{A}\right)\Gamma\left(k+\frac{b+A+r}{A}\right)}\\ &= \binom{k-1}{j}\left(\frac{b+A}{A}\right)_{j}\left(\frac{r}{A}\right)_{k-j}\left[\left(\frac{b+A+r}{A}\right)_k\right]^{-1}.\end{aligned} \tag{51}$$

Conditioning on $Z_0 = -v$, we get

$$\mathrm{P}\{N_{k-1} = k-1-j,\, Z_k = c \,|\, Z_0 = -v\} = \binom{k-1}{j}\left(\frac{r+A}{A}\right)_j\left(\frac{b}{A}\right)_{k-j}\left[\left(\frac{b+A+r}{A}\right)_k\right]^{-1} \tag{52}$$

and

$$\mathrm{P}\{N_{k-1}=k-1-j,\ Z_k=-v\,|\,Z_0=-v\}=\binom{k-1}{j}\left(\frac{r+A}{A}\right)_{j+1}\left(\frac{b}{A}\right)_{k-j-1}\left[\left(\frac{b+A+r}{A}\right)_k\right]^{-1}. \tag{53}$$

B Auxiliary functions

For the reader's convenience, now we provide some formulas used in the proofs.

The confluent hypergeometric function (or Kummer's function) used throughout the paper is so defined (see [1] for details)

$$ {}_1F_1(u,v;z)=\sum_{k=0}^{\infty}\frac{(u)_k}{(v)_k}\frac{z^k}{k!}\,. \tag{54}$$

We start recalling the relation

$$ {}_1F_1(a,b;z)=\mathrm{e}^{z}\,{}_1F_1(b-a,b;-z)\quad\text{or, equivalently,}\quad {}_1F_1(a,b;-z)=\mathrm{e}^{-z}\,{}_1F_1(b-a,b;z)\,. \tag{55}$$

An useful integration formula is Equation 3.383.1 of [15]:

$$\int_0^t y^{\nu-1}(t-y)^{\mu-1}\mathrm{e}^{\beta y}\mathrm{d}y=\frac{\Gamma(\mu)\Gamma(\nu)}{\Gamma(\mu+\nu)}t^{\mu+\nu-1}{}_1F_1(\nu,\mu+\nu;\beta t) \tag{56}$$

when $\mathrm{Re}(\mu)>0$ and $\mathrm{Re}(\nu)>0$.

Let X and Y be independent gamma-distributed random variables with parameters (α,μ) and (β,λ), respectively. By (55) and the integration formula (56), we find

$$F_X(t)=\frac{(\mu t)^{\alpha}}{\Gamma(\alpha+1)}\,\mathrm{e}^{-\mu t}\,{}_1F_1(1,\alpha+1;\mu t)\,,\qquad \mu>0,\ t\geq 0 \tag{57}$$

and so, for $\alpha,\ \beta,\ \mu,\ \lambda>0$ and $t\geq 0$,

$$\begin{aligned} G(\alpha,\mu,\beta,\lambda;t) &:= \mathrm{P}(X+Y\leq t)=\int_0^t F_X(t-y)f_Y(y)\mathrm{d}y \\ &=\mu^{\alpha}\lambda^{\beta}t^{\alpha+\beta}\mathrm{e}^{-\mu t}\sum_{h=0}^{+\infty}\frac{(\mu t)^h}{\Gamma(\alpha+\beta+h+1)}\,{}_1F_1(\beta,\alpha+\beta+h+1;(\mu-\lambda)t)\,. \end{aligned} \tag{58}$$

Note that an alternative expression of G was given in Moschopoulos [19] as series of suitable integrals. Analogously to (58), by (56) we can express the function

$$H\big(\alpha,\beta;t\big):=\int_0^t (t-y)\,{}_1F_1\big(1,2;\alpha(t-y)\big)\mathrm{e}^{-\alpha(t-y)}\mathrm{e}^{-\beta y}\,\mathrm{d}y\,,\qquad \alpha,\beta\in\mathbb{R},\ \ t\geq 0 \tag{59}$$

in terms of the hypergeometric function as

$$H\big(\alpha,\beta;t\big)=\begin{cases}\dfrac{t}{\alpha}\left[{}_1F_1(1,2;-\beta t)-\mathrm{e}^{-\alpha t}{}_1F_1(1,2;(\alpha-\beta)t)\right], & \alpha\neq 0,\ \beta\in\mathbb{R}\\[2mm] \dfrac{t}{\beta}\left[\mathrm{e}^{-\beta t}-\,{}_1F_1(1,2;-\beta t)\right], & \alpha=0,\ \beta\neq 0\\[2mm] \dfrac{t^2}{2}\,, & \alpha=0,\ \beta=0\,,\end{cases}$$

where we recall that ${}_1F_1(1,2;z)$ reduces to 1 when $z=0$ and to $(\mathrm{e}^{z}-1)/z$ when $z\neq 0$.

Acknowledgements.

This work was partially supported by MIUR (PRIN 2008) within the Projects "Mathematical models and computation methods for information processing and transmission in neuronal systems subject to stochastic dynamics" and "Bayesian methods: theoretical advancements and new applications".

References

[1] Abramowitz, M. and Stegun, I.A. (1994) *Handbook of Mathematical Functions with Formulas, Graph, and Mathematical Tables*. Dover, New York.

[2] Aletti, G., May, C. and Secchi, P. (2009) A central limit theorem, and related results, for a two-color randomly reinforced urn. *Adv. Appl. Prob.* **41**, 829-844.

[3] Bartlett, M.S. (1957) Some problems associated with random velocity. *Publ. Inst. Statist. Univ. Paris.* **6**, 261-270.

[4] Beghin, L., Nieddu, L. and Orsingher, E. (2001) Probabilistic analysis of the telegrapher's process with drift by means of relativistic transformations. *J. Appl. Math. Stochastic Anal.* **14**, 11-25.

[5] Berti, P., Crimaldi, I., Pratelli, L. and Rigo, P. (2011) A central limit theorem and its applications to multicolor randomly reinforced urns. *J. Appl. Prob.* **48**, 527-546.

[6] Berti, P., Pratelli, L. and Rigo, P. (2004) Limit theorems for a class of identically distributed random variables, *Ann. Prob.* **32**, 2029-2052.

[7] Crimaldi, I. (2009) An almost sure conditional convergence result and an application to a generalized Pólya urn, *Internat. Math. F.* **4**, no. 23, 1139-1156.

[8] De Gregorio, A. (2010) Stochastic velocity motions and processes with random time. *Adv. Appl. Prob.* **42**, 1028-1056.

[9] Di Crescenzo, A. (2001) On random motions with velocities alternating at Erlang-distributed random times. *Adv. Appl. Prob.* **33**, 690-701.

[10] Di Crescenzo, A. and Martinucci, B. (2007) Random motion with gamma-distributed alternating velocities in biological modeling. In *Computer Aided Systems Theory, EUROCAST 2007*; Moreno-Díaz, R. *et al.*, Eds.; *Lect. Notes Comp. Science* **4739**, 163-170. Springer-Verlag, Berlin.

[11] Di Crescenzo, A. and Martinucci, B. (2010) A damped telegraph random process with logistic stationary distribution. *J. Appl. Prob.* **47**, 84-96.

[12] Di Crescenzo, A., Martinucci, B. and Zacks, S. (2011) On the damped geometric telegrapher's process. In *Mathematical and Statistical Methods for Actuarial Sciences and Finance*; Perna, C., Sibillo, M., Eds.; p. 175-182. Springer, Dordrecht.

[13] Foong, S.K. and Kanno, S. (1994) Properties of the telegrapher's random process with or without a trap. *Stoch. Proc. Appl.* **53**, 147-173.

[14] Goldstein, S. (1951) On diffusion by discontinuous movements, and on the telegraph equation. *Quart. J. Mech. Appl. Math.* **4**, 129-156.

[15] Gradshteyn, I.S. and Ryzhik, I.M. (2007) *Tables of Integrals, Series and Products*, 7th Ed. Academic Press, Amsterdam.

[16] Iacus, S.M. (2001) Statistical analysis of the inhomogeneous telegrapher's process. *Stat. Prob. Lett.* **55**, 83-88.

[17] Kac, M. (1974) A stochastic model related to the telegrapher's equation. *Rochy Mount. J. Math.* **4**, 497-509.

[18] Mahmoud, H.M. (2008) *Pólya Urn Models*, CRC press, Taylor & Francis.

[19] Moschopoulos, P.G. (1985) The distribution of the sum of independent gamma random variables. *Ann. Inst. Statist. Math.* **37**, 541-544.

[20] Orsingher, E. (1990) Probability law, flow functions, maximum distribution of wave-governed random motions and their connections with Kirchoff's laws. *Stoch. Process. Appl.* **34**, 49-66.

[21] Orsingher, E. (1995) Motions with reflecting and absorbing barriers driven by the telegraph equation. *Random Oper. Stoch. Equat.* **1**, 9-21.

[22] Pinsky, M.A. (1991) *Lectures on Random Evolutions*. World Scientific, Singapore.

[23] Pólya, G. (1931) Sur quelques points de la théorie des probabilités. *Ann. Inst. Poincaré* **1**, 117-161.

[24] Stadje, W. and Zacks, S. (2004) Telegraph processes with random velocities. *J. Appl. Prob.* **41**, 665-678.

[25] Wolff, R.W. (1989) *Stochastic modeling and the theory of queues*. Prentice Hall International Series in Industrial and Systems Engineering. Prentice Hall, Inc., Englewood Cliffs, NJ.

[26] Zacks, S. (2004) Generalized integrated telegraph processes and the distribution of related random times. *J. Appl. Prob.* **41**, 497-507.